

**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**

**Wound Care:**

- Remove dressing in 2 days and apply band aid to cover sutures.
- Keep steri-strips in place until they fall off or they are removed at your first post-operative visit.
- Showering after 4 days is fine; be sure to pat the incision dry afterwards.
- Do not scrub the area; just allow water/soap to wash over you.
- Do not bath or swim until the sutures are removed.
- Continue to wear your TEDS stockings for one week after surgery to help decrease the swelling. You may remove for showering, etc.

**Ice:**

- We recommend that you use the ice wrap/cryocuff on a consistent basis for the first 24–48 hours. This will help reduce post-operative swelling. After that, use as necessary.
- If the cryocuff is not available or if you find this uncomfortable, you can simply apply an ice pack to your knee 3 – 4 times per day until the swelling resolves.

**Braces/Devices:**

- With an **Anterior Cruciate Ligament (ACL) Reconstruction with Meniscal Repair**, you will wear your Bledsoe Brace in the locked position when sleeping and walking (your leg will be fully extended). Otherwise, it should be set from 0-90 degrees for 4 weeks, followed by full range of motion. After 4 weeks, you may bear weight with the knee brace unlocked. You need to wear the brace for a total of 6 weeks.
- *If you have a question about your brace or you need help adjusting it, please call NOPCO at (617) 355.6887.*
- You have been given a CPM (motion machine). You may use it without your brace. It should be set from 0-110 degrees. You should use the CPM 2-3 times a day for 2-3 hrs at a time.
- *If you have a question about your CPM and its settings, please call MED at (781) 337.3070.*

**Crutches & Weight bearing status:**

- Following an **Anterior Cruciate Ligament (ACL) Reconstruction with Meniscal Repair**, you will be able to partially bear weight (touch the floor lightly with your toes for balance) with crutches for the first 2 weeks if you have had an autograft repair (use of your own tendon), and for 4 weeks with an allograft repair (use of a donated tendon). Dr. Ramappa will tell you which procedure you have had.


**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**

**Physical Therapy:**

- Formal physical therapy will be ordered by your orthopedic surgeon in the form of a written prescription upon discharge from surgery. You will also receive a therapy protocol form. Following an **anterior cruciate ligament (ACL) reconstruction with Meniscal Repair**, you should start formal physical therapy within 2 - 4 days. The prescription and protocol should be given to a physical therapist of your choice so you can begin your program as directed under close supervision of your surgeon.
  
- Until you begin working with a therapist, you can do some exercises at home, such as: quad sets, leg raises and calf pumps (see pictures below).

**Quad Set Exercise**

- a. Tighten the muscles on top of the thigh as tightly as possible and hold.
  1. Pull your toes back.
  2. Push the back of your knee down to the floor.
  3. Try to push out and up through the heel.
- b. Pull 10 seconds, trying every second to pull even tighter.
- c. Relax 5 seconds.
- d. Repeat for 2 sets of ten times. Rest 60 seconds between sets.



**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**


**Straight Leg Raise**

- a. Tighten the muscles on top of the thigh as tightly as possible and hold.
- b. Raise the entire leg holding the knee as tight as possible. Hold 5 seconds.
- c. Lower leg and rest 2 seconds.
- d. Repeat for 2 sets of 10 times.
- e. Rest 1 minute between sets.



**Flexion and Extension**

- a. Sit on something high enough to keep foot off the floor.
- b. Bend the knee as far back as possible.
- c. Straighten knee as far forward as possible and hold it straight.
- d. Then relax. (Like pumping on a swing.)
- e. Repeat for 3 sets of 10 times.



**Calf Pumps**

**Pointing the Feet:- Action**

1. (Keeping your foot strictly in line with the ankle knee and hip joints): Point the foot away from you.
2. Repeat slowly, five to 10 times each foot.

**Flexing Feet:- Action**

1. (Keeping your foot strictly in line with the ankle knee and hip joints): Flex the foot, this time letting the heel push away from you, and the toe end of the foot come toward you.
2. Repeat slowly, five to 10 times each foot.

Flexing the Foot      Pointing the Foot



**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**



**Pain Control:**

***Narcotics***

- You have been prescribed \_\_\_\_\_, which is a narcotic pain medication.
- Use the medication only as prescribed. Take it with food.
- Do not drive or use any type of heavy machinery, drink alcohol, make any life-changing or legal decisions (i.e. sign a will), or participate in activities that require a lot of physical skill.
- Take a stool softener, such as Colace, while taking the narcotic medication, because narcotics may cause constipation.

***Non- Steroidal Anti-Inflammatory Medications:***

- Please take an over the counter non-steroidal anti-inflammatory such as Motrin, Advil, Ibuprofen (600 mg every 12 hours) or Aleve (500 mg every 12 hours) for additional pain control, only if you do not have a contraindication to taking this medication (i.e. allergy to the medication, stomach ulcers or kidney problems).
- Take this medication with food, and try to stagger the times you take this medication with your pain medication. This may help decrease the chance of stomach upset.
- If you can take Aspirin without any difficulty, then you should take **Aspirin 325 mg daily for 4 weeks, starting the day of your surgery, upon arriving home.**

**Driving:**

**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**

- In general, you can resume driving after your first post-op appointment, when you have full control of your arms and legs and you are pain free. Do not drive if you are still taking narcotic pain medications and/or you are still partial weight bearing.

**Return to Work:**

- You may return to work as soon as you are comfortable and able to safely weight bear using crutches. This typically occurs 2–4 weeks post-op. Return to work notes can be obtained from your orthopedic surgeon at your first post-op appointment.

**Follow-up Appointment:**

- Your first post-operative appointment will be scheduled 10 – 14 days following your surgical procedure. At this appointment your sutures will be removed. If you do not have a post-operative appointment scheduled when you leave following surgery, please call 617-667-3940 to make the appointment. Your first post-operative appointment will be with a Physicians' Assistant (PA) or a Nurse Practitioner (NP).

**\*\*\*\*Signs & Symptoms to Immediately Report\*\*\*\***

- ◆ *Call 911 and go to the nearest hospital if you are having chest pain or trouble breathing.*

**Call the office at: 617-667-3940 to report any of the following:**

- ◆ Persistent fever (101 or greater)
- ◆ Sudden increase in pain and swelling
- ◆ Wound redness or drainage
- ◆ Increased skin temperature around incision
- ◆ Deep calf pain and swelling

**DR. RAMAPPA'S POST-OPERATIVE INSTRUCTIONS**  
**Anterior Cruciate Ligament (ACL) Reconstruction with MENISCAL REPAIR**  
**Sports Medicine**  
**Beth Israel Deaconess Medical Center**  
**617-667-3940**

DATE: 4/27/08

Sources:

Huddleston, HD, Dr.; The Hip and Knee Institute; Home Exercises for the First 8 Weeks.  
Pages 1-4.